ADDENDUM TO "ON SUPER-JORDANIAN $\mathcal{U}_{\mathsf{h}}(sl(N|1))$ ALGEBRA"

B. ABDESSELAM[a], A. CHAKRABARTI[b], R. CHAKRABARTI[c], A. YANALLAH[d] and M.B ZAHAF[e]

[a,d,e]*Laboratoire de Physique Quantique de la Matière et Modélisations Mathématiques (LPQ3M), Centre Universitaire de Mascara, 29000-Mascara, Algérie*

[a]*Laboratoire de Physique Théorique d'Oran, Université d'Oran Es-Sénia, 31100-Oran, Algérie*

[b]*Centre de Physique Théorique, Ecole Polytechnique, 91128-Palaiseau cedex, France.*

[c]*Department of Theoretical Physics, University of Madras, Guindy Campus, Madras 600025, India*

arXiv:math/0701379v1 [math.QA] 13 Jan 2007

Abstract

We give a complete proof of the result (2.10) presented in our paper published in **J. Phys. A: Math. Gen. 39 (2006) 8307–8319**.

1. Let $\mathcal{U}_q(sl(2|1))$ (q is an arbitrary complex number) be the Hopf superalgebra generated by the elements $\hat{h}_i$, $\hat{e}_i$ and $\hat{f}_i$, $i = 1, 2$, satisfy the relations

$$\begin{aligned} &[\hat{h}_i,\ \hat{h}_j] = 0, \qquad [\hat{h}_i,\ \hat{e}_j] = a_{ij}\hat{e}_j, \qquad [\hat{h}_i,\ \hat{f}_j] = -a_{ij}\hat{f}_j, \\ &[\hat{e}_i,\ \hat{f}_j] = \delta_{ij}\frac{q^{\hat{h}_i} - q^{-\hat{h}_i}}{q - q^{-1}}, \qquad \hat{e}_2^2 = \hat{f}_2^2 = 0, \\ &\hat{e}_1^2\hat{e}_2 - \left(q + q^{-1}\right)\hat{e}_1\hat{e}_2\hat{e}_1 + \hat{e}_2\hat{e}_1^2 = 0, \qquad \hat{f}_1^2\hat{f}_2 - \left(q + q^{-1}\right)\hat{f}_1\hat{f}_2\hat{f}_1 + \hat{f}_2\hat{f}_1^2 = 0, \end{aligned} \tag{1}$$

where $[\ ,\ ]$ is the supercommutator given by $[a,\ b] = ab - (-)^{\deg(a)\deg(b)}ba$ and $a_{11} = 2$, $a_{12} = a_{21} = -1$, $a_{22} = 0$. All generators are even except for $\hat{e}_2$ and $\hat{f}_2$ which are odd. The coproducts, counits and antipodes are given by

$$\begin{aligned} &\Delta\left(\hat{e}_i\right) = \hat{e}_i \otimes q^{\hat{h}_i/2} + q^{-\hat{h}_i/2} \otimes \hat{e}_i, \qquad \epsilon(\hat{e}_i) = 0, \qquad S(\hat{e}_i) = -q^{\hat{h}_i/2}\hat{e}_i q^{-\hat{h}_i/2}, \\ &\Delta\left(\hat{f}_i\right) = \hat{f}_i \otimes q^{\hat{h}_i/2} + q^{-\hat{h}_i/2} \otimes \hat{f}_i, \qquad \epsilon(\hat{f}_i) = 0, \qquad S(\hat{f}_i) = -q^{\hat{h}_i/2}\hat{f}_i q^{-\hat{h}_i/2}, \\ &\Delta\left(\hat{h}_i\right) = \hat{h}_i \otimes 1 + 1 \otimes \hat{h}_i, \qquad \epsilon(\hat{h}_i) = 0, \qquad S(\hat{h}_i) = -\hat{h}_i. \end{aligned} \tag{2}$$

Let $\hat{e}_3 = \hat{e}_1\hat{e}_2 - q^{-1}\hat{e}_2\hat{e}_1$ and $\hat{f}_3 = \hat{f}_2\hat{f}_1 - q\hat{f}_1\hat{f}_2$. Khroshkin, Tolstoy [1] and Yamane [2] showed that

$$\mathcal{R}_q = \hat{\mathcal{R}}_q K_q, \tag{3}$$

where

$$\begin{aligned} K_q &= q^{-h_1\otimes h_2 - h_2\otimes h_1 - 2h_2\otimes h_2}, \\ \hat{\mathcal{R}}_q &= \exp_q\left[\left(q - q^{-1}\right)q^{-h_1/2}\hat{e}_1 \otimes \hat{f}_1 q^{h_1/2}\right]\exp_q\left[-\left(q - q^{-1}\right)q^{-h_1/2-h_2/2}e_3 \otimes \hat{f}_3 q^{h_1/2+h_2/2}\right] \times \\ &\quad \exp_q\left[-\left(q - q^{-1}\right)q^{-h_2/2}\hat{e}_2 \otimes \hat{f}_2 q^{h_2/2}\right] \end{aligned} \tag{4}$$

and $\exp_q(x) = \sum_{n\geq 0} x^n/(n)_q!$, $(n)_q! = (1)_q(2)_q\cdots(n)_q$, $(n)_q = \frac{1-q^k}{1-q}$. In the (fund.)⊗(arbitrary) representation, the R-matrix (3) takes the following form:

$$R_{\mathsf{q}}\big|_{(fund.\otimes arb.)} = \begin{pmatrix} q^{-h_2} & A & B \\ 0 & q^{-h_1-h_2} & C \\ 0 & 0 & q^{-h_1-2h_2} \end{pmatrix}, \tag{5}$$

where

$$\begin{aligned} A &= \left(q - q^{-1}\right)q^{-1/2}\hat{f}_1 q^{-h_1/2-h_2}, \\ B &= -\left(q - q^{-1}\right)^2 q^{-1}\hat{f}_1 q^{h_1/2}\hat{f}_2 q^{-h_1-3h_2/2} - \left(q - q^{-1}\right)q^{-1/2}\hat{f}_3 q^{-h_1/2-3h_2/2}, \\ C &= \left(q - q^{-1}\right)q^{-1/2}\hat{f}_2 q^{-h_1-3h_2/2}, \end{aligned} \tag{6}$$

2. The R_{h}-matrix in the (fund.)⊗(arbitrary) representation is obtained, from (5), as follows:

$$\begin{aligned} R_{\mathsf{h}}\big|_{(fund.\otimes arb.)} &= \lim_{q\to 1}\begin{pmatrix} \mathsf{G}^{-1} & -\frac{\mathsf{h}}{q-1}\mathsf{G}^{-1} & 0 \\ 0 & \mathsf{G}^{-1} & 0 \\ 0 & 0 & \mathsf{G}^{-1} \end{pmatrix} R_{\mathsf{q}}\big|_{(fund.\otimes arb.)} \begin{pmatrix} \mathsf{G} & \frac{\mathsf{h}}{q-1}\mathsf{G} & 0 \\ 0 & \mathsf{G} & 0 \\ 0 & 0 & \mathsf{G} \end{pmatrix} \\ &= \lim_{q\to 1}\begin{pmatrix} \mathsf{G}^{-1}q^{-h_2}\mathsf{G} & \alpha & \beta \\ 0 & \mathsf{G}^{-1}q^{-h_1-h_2}\mathsf{G} & \gamma \\ 0 & 0 & \mathsf{G}^{-1}q^{-h_1-2h_2} \end{pmatrix}, \end{aligned} \tag{7}$$

where

$$\begin{aligned} \alpha &= \frac{\mathsf{h}}{q-1}\left(\mathsf{G}^{-1}q^{-h_2}\mathsf{G} - \mathsf{G}^{-1}q^{-h_1-h_2}\mathsf{G}\right) + \left(q-q^{-1}\right)q^{-1/2}\mathsf{G}^{-1}\hat{f}_1 q^{-h_1/2-h_2}\mathsf{G}, \\ \beta &= -\left(q-q^{-1}\right)^2 q^{-1}\mathsf{G}^{-1}\hat{f}_1 q^{h_1/2}\hat{f}_2 q^{-h_1-3h_2/2}\mathsf{G} - \left(q-q^{-1}\right)q^{-1/2}\mathsf{G}^{-1}\hat{f}_3 q^{-h_1/2-3h_2/2}\mathsf{G} + \\ &\quad \frac{\mathsf{h}}{q-1}\left(q-q^{-1}\right)q^{-1/2}\mathsf{G}^{-1}\hat{f}_2 q^{-h_1-3h_2/2}\mathsf{G} \\ \gamma &= -\left(q-q^{-1}\right)q^{-1/2}\mathsf{G}^{-1}\hat{f}_2 q^{-h_1-3h_2/2}\mathsf{G} \end{aligned} \tag{8}$$

and $\mathsf{G} = E_q\left(\frac{\mathsf{h}}{q-1}\hat{e}_1\right)$, $E_q(x) = \sum_{n\geq 0} x^n/[n]!$, $[n]! = [n]\cdots[1]$, $[n] = \frac{q^n-q^{-n}}{q-q^{-n}}$.

3. Defining

$$t^{(\alpha)} = E_q^{-1}\left(\frac{\mathsf{h}}{q-1}\hat{e}_1\right)E_q\left(q^{\alpha}\frac{\mathsf{h}}{q-1}\hat{e}_1\right), \qquad t^{(0)} = 1, \tag{9}$$

we obtain the following properties:

$$E_q^{-1}\left(\frac{\mathsf{h}\hat{e}_1}{q-1}\right)q^{\alpha h_1/2}E_q\left(\frac{\mathsf{h}\hat{e}_1}{q-1}\right) = t^{(\alpha)}q^{\alpha h_1/2}, \tag{10}$$

$$t^{(\alpha+\beta)}q^{(\alpha+\beta)h_1/2} = t^{(\alpha)}q^{\alpha h_1/2}t^{(\beta)}q^{\beta h_1/2}, \tag{11}$$

$$E_q^{-1}\left(\frac{\mathsf{h}\hat{e}_1}{q-1}\right)q^{\beta h_2}E_q\left(\frac{\mathsf{h}\hat{e}_1}{q-1}\right) = t^{(-\beta)}q^{\beta h_2}, \tag{12}$$

$$E_q^{-1}\left(\frac{\mathsf{h}\hat{e}_1}{q-1}\right)\hat{f}_1 E_q\left(\frac{\mathsf{h}\hat{e}_1}{q-1}\right) = \hat{f}_1 - \frac{\mathsf{h}}{(q-1)\left(q-q^{-1}\right)}\left(t^{(1)}q^{h_1} - t^{(-1)}q^{-h_1}\right), \tag{13}$$

$$E_q^{-1}\left(\frac{\mathsf{h}\hat{e}_1}{q-1}\right)\hat{f}_2 E_q\left(\frac{\mathsf{h}\hat{e}_1}{q-1}\right) = \hat{f}_2, \tag{14}$$

$$E_q^{-1}\left(\frac{\mathsf{h}\hat{e}_1}{q-1}\right)\hat{f}_3 E_q\left(\frac{\mathsf{h}\hat{e}_1}{q-1}\right) = \hat{f}_3 + \frac{\mathsf{h}q}{q-1}t^{(1)}\hat{f}_2 q^{h_1}. \tag{15}$$

4. Let us introduce the following generator:

$$T = \lim_{q\to 1} t^{(1)}. \tag{16}$$

From (11) it is evident that

$$\lim_{q\to 1} t^{(\alpha)} = T^{\alpha}. \tag{17}$$

To obtain a closed form of T, we proceed as follows: we left and right multiply the commutation relation $q^{h_1} - q^{-h_1} = \left(q-q^{-1}\right)\left(\hat{e}_1\hat{f}_1 - \hat{f}_1\hat{e}_1\right)$ by G^{-1} and G respectively. After simple calculations, we reach to

$$t^{(2)}q^{h_1} - t^{(-2)}q^{-h_1} = q^{h_1} - q^{-h_1} + \mathsf{h}(q+1)\left[t^{(1)}\hat{e}_1 q^{h_1} + q^{-h_1}\hat{e}_1 t^{(-1)}\right], \tag{18}$$

which yields, when $q \longrightarrow 1$, to

$$T^2 - T^{-2} = 2\mathsf{h}\left(T + T^{-1}\right)\hat{e}_1 \qquad \Rightarrow \qquad T - T^{-1} = 2\mathsf{h}\hat{e}_1. \tag{19}$$

Finally, we obtain

$$T^{\pm 1} = \pm \mathsf{h}\hat{e}_1 + \sqrt{1 + \mathsf{h}^2 \hat{e}_1^2}. \tag{20}$$

5. We turn now to (7). It is easy to verify that

$$\begin{aligned}
&\lim_{q\to 1} \mathsf{G}^{-1} q^{-h_2} \mathsf{G} = \lim_{q\to 1} t^{(1)} q^{-h_2} = T, \\
&\lim_{q\to 1} \mathsf{G}^{-1} q^{-h_1-h_2} \mathsf{G} = \lim_{q\to 1} t^{(-1)} q^{-h_1-h_2} = T^{-1}, \\
&\lim_{q\to 1} \mathsf{G}^{-1} q^{-h_1-2h_2} \mathsf{G} = 1, \\
&\lim_{q\to 1} \gamma = -\lim_{q\to 1} \left(q - q^{-1}\right) q^{-1/2} \hat{f}_2 t^{(-1/2)} q^{-h_1-3h_2/2} = 0, \\
&\lim_{q\to 1} \alpha = \lim_{q\to 1} \left(q - q^{-1}\right) q^{-1/2} \hat{f}_1 + \frac{\mathsf{h} t^{(1)}}{q-1} \left(q^{-h_2} - q^{-h_1/2-h_2-1/2}\right) - \frac{\mathsf{h} t^{(-1)}}{q-1} \left(q^{-h_1-h_2} - q^{-3h_1/2-h_2-1/2}\right) \\
&\qquad = -\frac{\mathsf{h}}{2} \left(T + T^{-1}\right) h_1 + \frac{\mathsf{h}}{2} \left(T - T^{-1}\right) \\
&\qquad \equiv -\mathsf{h} H_1 + \frac{\mathsf{h}}{2} \left(T - T^{-1}\right), \\
&\lim_{q\to 1} \beta = -\lim_{q\to 1} \left(q - q^{-1}\right)^2 q^{-1} \left[\hat{f}_1 - \frac{\mathsf{h}}{(q-1)\left(q - q^{-1}\right)} \left(t^{(1)} q^{h_1} - t^{(-1)} q^{-h_1}\right)\right] t^{(1)} q^{h_1^{/}2} \hat{f}_2 t^{(-1/2)} q^{-h_1-3h_2/2} \\
&\qquad\quad - \left(q - q^{-1}\right) q^{-1/2} \left[f_3 + \frac{\mathsf{h} q}{q-1} t^{(1)} q^{h_1}\right] t^{(1/2)} q^{-h_1/2-3h_2/2} \\
&\qquad\quad + \frac{\mathsf{h}\left(q - q^{-1}\right)}{q-1} q^{-1/2} \hat{f}_2 t^{(-1/2)} q^{-h_1-3h_2/2} \\
&\qquad = 2\mathsf{h} \left(T - T^{-1}\right) T^{1/2} f_2 - 2\mathsf{h} T^{3/2} f_2 + 2\mathsf{h} T^{-1/2} f_2 = 0.
\end{aligned} \tag{21}$$

Finally, we obtain

$$R_{\mathsf{h}}\big|_{(fund.\otimes fund.)} = \begin{pmatrix} T & -\mathsf{h} H_1 + \frac{\mathsf{h}}{2}\left(T - T^{-1}\right) & 0 \\ 0 & T^{-1} & 0 \\ 0 & 0 & 1 \end{pmatrix}. \tag{22}$$

References

[1] Khoroshkin S.M. and Tolstoy V.N., Comm. Math. Phys. **141** (1991), 599.

[2] Yamane H., Publ. Res. Inst. Math. Sci. **30** (1994), 1587.